

KIDNEY AND PANCREAS TRANSPLANT PROGRAM

Living Kidney Donation Selection/Exclusion Criteria

Absolute Contraindications to Living Kidney Donation

- Patient both under the age of 18 years and mentally incapable of making an informed decision
- HIV positive
- Active malignancy, or incompletely treated malignancy that either:
 - Requires treatment other than surveillance or
 - Has more than minimal known risk transmission
- Diabetes Mellitus; Gestational Diabetes
- High suspicion of donor inducement, coercion, or other undue pressure
- High suspicion of knowingly and unlawfully acquiring, receiving, or otherwise transferring anything of value in exchange for any human organ
- Evidence of acute symptomatic infection (until resolved)
- Uncontrolled diagnosable psychiatric conditions requiring treatment before donation, including any evidence of suicidality
- Uncontrollable hypertension or history of hypertension with evidence of end organ damage
- Current pregnancy
- History of pre-eclampsia in women

Relative Contraindications to Living Kidney Donation

- Abnormal urinalysis results
 - Proteinuria
 - Microscopic Hematuria
 - Microscopic Pyuria
 - Abnormal glomerular filtration rate (GFR)
- Abnormal Renal Imaging by CT Angiogram
 - Abnormal Parenchyma
 - Abnormal Vasculature
- Nephrolithiasis
- Hypertension
- Hereditary Renal Disease
 - Autosomal dominant polycystic kidney disease
 - Alport Syndrome
 - TBM Disease (benign familial hematuria)
- Systemic Lupus Erythematosus (SLE)
- Familial Primary Glomerulonephritis
- Body Mass Index (BMI) >32 kg/m²
- Mental disorders
- Substance Abuse



AFFILIATED WITH MEDICAL COLLEGE OF GEORGIA